

NO-1192 877

MEMORY-INDUCED EXTERIOR RESONANCES OF MOSCOWITES(0) STATE 2/1

UNIV OF NEW YORK AT BUFFALO DEPT OF CHEMISTRY

H F ARNOLDUS ET AL. MAR 88 UBUFFALO/DC/88/TR-66


NO0014-86-K-0043

F/G 20/5

NL

UNCLASSIFIED





MICROCOPY RESOLUTION TEST CHART  
 (1963-A)

AD-A192 877

OFFICE OF NAVAL RESEARCH

Contract N00014-86-K-0043

TECHNICAL REPORT No. 66

Memory-Induced Extra Resonances of Adsorbates

by

Henk F. Arnoldus and Thomas F. George

Prepared for Publication

in

Physical Review Letters

Departments of Chemistry and Physics  
State University of New York at Buffalo  
Buffalo, New York 14260

March 1988

Reproduction in whole or in part is permitted for any purpose of the  
United States Government.

This document has been approved for public release and sale;  
its distribution is unlimited.

DTIC  
ELECTE  
MAR 15 1988  
S H D

139

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

## REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) UBUFFALO/DC/88/TR-66			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Depts. Chemistry & Physics State University of New York		6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION		
6c. ADDRESS (City, State, and ZIP Code) Fronczak Hall, Amherst Campus Buffalo, New York 14260			7b. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Contract N00014-86-K-0043		
8c. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.
			WORK UNIT ACCESSION NO.		
11. TITLE (Include Security Classification) Memory-Induced Extra Resonances of Adsorbates					
12. PERSONAL AUTHOR(S) Henk F. Arnoldus and Thomas F. George					
13a. TYPE OF REPORT		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) March 1988	
15. PAGE COUNT 11					
16. SUPPLEMENTARY NOTATION Prepared for publication in Physical Review Letters (in press)					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	ADSORBATES; EXTRA RESONANCES; MEMORY-INDUCED		
			OPTICAL ABSORPTION PROFILE MARKOVIAN LIMIT MODIFIED LORENTZIAN		
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
<p>The optical absorption profile of an atom-surface vibrational bond is studied. From Markovian relaxation theory it follows that the line shape is a Lorentzian around the adsorbate resonance frequency <math>\omega_0</math>. Dispersion relations for crystals have a fairly small cutoff frequency <math>\omega_D</math>, which prohibits the use of such a simple theory. We calculated the spectral profile with finite memory-time reservoir theory, and we found that the modified Lorentzian vanishes above <math>\omega_D</math>. Also a new spectral line at <math>\omega_0 + \omega_D</math> is predicted, which disappears in the Markovian limit. The physical origin of the new line is explained.</p>					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nelson			22b. TELEPHONE (Include Area Code) (202) 696-4410		22c. OFFICE SYMBOL

DD Form 1473, JUN 86

Previous editions are obsolete.

SECURITY CLASSIFICATION OF THIS PAGE

UNCLASSIFIED


MEMORY-INDUCED EXTRA RESONANCES OF ADSORBATES

Henk F. Arnoldus and Thomas F. George  
Departments of Physics and Chemistry  
239 Fronczak Hall  
State University of New York at Buffalo  
Buffalo, New York 14260

ABSTRACT

The optical absorption profile of an atom-surface vibrational bond is studied. From Markovian relaxation theory it follows that the line shape is a Lorentzian around the adsorbate resonance frequency  $\omega_0$ . Dispersion relations for crystals have a fairly small cutoff frequency  $\omega_D$ , which prohibits the use of such a simple theory. We calculated the spectral profile with finite memory-time reservoir theory, and we found that the modified Lorentzian vanishes above  $\omega_D$ . Also a new spectral line at  $\omega_0 + \omega_D$  is predicted, which disappears in the Markovian limit. The physical origin of the new line is explained.

PACS: 32.70.-n, 32.80.-t, 71.36.+c, 78.90.+t



Accession For	
NTIS CPA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

Adsorbed atoms on the surface of a crystal can absorb photons from an incident infrared laser beam. Coupling between the van der Waals bond and the radiation is brought about by the motion-induced dipole moment  $\mu$  (the atom itself is assumed to be neutral for IR light), and the atom-crystal interaction is governed by the single-phonon coupling Hamiltonian<sup>1</sup>

$$H_i = -(\underline{u} \cdot \underline{e}_z) dV/dz, \quad (1)$$

in terms of the derivative of the binding potential well  $V(z)$ . Here  $z$  denotes the normal direction to the surface (we neglect lateral motion) and  $\underline{u}$  is the phonon-field amplitude operator,<sup>2</sup> evaluated at the position of the surface atom which is closest to the adsorbate. Relaxation of the atomic bond due to phonon exchange with the crystal is most conveniently described with reservoir theory, where the crystal is regarded as a thermal bath.<sup>3</sup> Then an equation of motion for the reduced adsorbate density operator can be derived, in which the properties of the heat bath only enter parametrically, and can be expressed in terms of the Fourier-Laplace transform of the reservoir correlation function

$$g(\omega/\omega_D) = \frac{2M\omega_D^2}{3\pi\hbar} \int_0^\infty d\tau e^{i\omega\tau} \langle (\underline{u}(\tau) \cdot \underline{e}_z) (\underline{u} \cdot \underline{e}_z) \rangle. \quad (2)$$

Equation (2) defines the dimensionless function  $g(z)$  in terms of the mass  $M$  of a bulk atom and the Debye frequency  $\omega_D$ . Then it can be shown that  $g(z)$  depends only on the dimensionless temperature  $\gamma = \hbar\omega_D/k_B T$ , with  $k_B$  Boltzmann's constant. An explicit expression for  $g(z)$  can be found in Ref. 4.

In the most simple theory of relaxation<sup>5</sup> one adopts the Markov and secular approximations, which yields for the low-intensity absorption line shape as a function of the laser frequency  $\omega$

$$I(\omega) = \omega_0 (n_1 - n_2) \text{Re} \frac{i}{\omega - \omega_0 + ia(g(\hat{\omega}_0) + g^*(-\hat{\omega}_0))} \quad (3)$$

This profile is a Lorentzian with its peak value at  $\omega = \omega_0 + a\text{Im}(g(\hat{\omega}_0) + g^*(-\hat{\omega}_0))$ , and a half-width at half-maximum equal to  $a\text{Re}(g(\hat{\omega}_0) + g^*(-\hat{\omega}_0))$ . The notation is  $\hat{\omega} = \omega/\omega_D$ ,  $\omega_0$  = unperturbed adsorbate resonance, and  $n_i$  = population of the  $i$ -th level ( $n_1 > n_2$ ). The temperature dependence is incorporated in  $g(z)$ , and the phonon-coupling strength is measured by

$$a = \frac{3\pi}{2M\omega_D^2} \langle 2 \left| \frac{dV}{dz} \right| 1 \rangle^2 \quad (4)$$

Most crucial for the derivation of the line profile (3) is the conjecture that the phonon-amplitude correlation function  $\langle (u(\tau) \cdot e_z)(u \cdot e_z) \rangle$  decays to zero very fast for  $\tau > 0$ , which would justify the Markov approximation. It is an essential feature of a crystal, however, that its dispersion relation has a finite cutoff frequency  $\omega_D$ , which sets the lower limit of  $1/\omega_D$  on the decay time. In fact, the structure of the amplitude operator  $u$  is such that the correlation function decays only as  $1/\tau$  for  $\tau \rightarrow \infty$ , rather than exponentially, which makes a Markov approximation (zero memory time) at least doubtful. In order to investigate this problem quantitatively, we have developed a finite memory-time relaxation theory,<sup>6</sup> with no restriction on the time scales. Subsequently, we have applied this theory to the evaluation of the absorption line shape, with the formal result<sup>7</sup>

$$I(\omega) = \omega \mu_{21}^{-2} \text{ReTr} L_X \frac{i}{\omega - L_a + i\Gamma(\omega)} (L_Y - i\Gamma(\omega)) \bar{\rho} \quad (5)$$

This expression holds for any configuration of adsorbate levels and includes multiphonon transitions as well. In Eq. (5),  $\bar{\rho}$  equals the adsorbate density operator in thermal equilibrium, and the Liouvillians  $L_X$  and  $L_Y$  are defined by  $L_X \sigma = \mu \sigma$ ,  $L_Y \sigma = [\mu, \sigma]$  in terms of their action on an arbitrary adsorbate density operator  $\sigma$ . The relaxation operator  $\Gamma(\omega)$  is frequency dependent, which reflects the memory in the time regression of two-time quantum correlation functions. In the time domain this would be a memory kernel,<sup>8</sup> and in the Markov approximation  $\Gamma(\omega)$  assumes an  $\omega$ -independent value. Furthermore, a Liouville operator  $T(\omega)$  appears, which accounts for the fact that the density operator of the entire system does not factorize as  $\rho_{\text{adsorbate}} \times \rho_{\text{crystal}}$  in thermal equilibrium if the memory time is finite. We call this term the initial correlation contribution to the line shape, in contrast to the term  $L_Y$ , which is referred to as the regression part. In a zero memory-time approximation the operator  $T(\omega)$  vanishes identically, since then a factorization of the density operator can be justified. Both  $\Gamma(\omega)$  and  $T(\omega)$  are complicated operators. Elsewhere<sup>7</sup> we have studied the modifications of the Lorentzian profile (3) due to the memory effects, and we have derived an explicit expression for  $I(\omega)$ , pertaining to the situation where  $\omega_0$  is smaller than  $\omega_D$ .

In this Letter we report a peculiar feature of the absorption profile (5) which arises as a consequence of the memory in the interaction. Since Eq. (5) applies to any configuration, we can consider the case where the vibrational resonance  $\omega_0$  is larger than  $\omega_D$ . Then we scan the laser over this resonance, implying that also  $\omega > \omega_D$ . For frequencies larger than  $\omega_D$  a Markov approximation can never be justified, and it is imperative to include the memory in the time evolution. At this stage we neglect the small contribution of multiphonon



processes, in order to emphasize the significance of the predicted effects. Then it can be shown<sup>9</sup> that the regression part ( $\sim L_Y$ ), which would lead to the Lorentzian in the Markov approximation, is identically zero for  $\omega > \omega_D$ . Hence any absorption is due to the initial-correlation term  $T(\omega)$ , and therefore a mere consequence of the finiteness of the memory time. We obtain for the absorption spectrum

$$I(\omega) = a \left( \frac{\mu_{22} - \mu_{11}}{\mu_{21}} \right)^2 \operatorname{Re} \frac{g(\hat{\omega} - \hat{\omega}_0)}{\omega + ia(g(\hat{\omega} - \hat{\omega}_0) + g^*(\hat{\omega}_0 - \hat{\omega}))} \quad (6)$$

for  $\omega > \omega_D$ . Here we used that the population of the upper state must be zero for  $\omega_0 > \omega_D$  (so  $n_1 = 1$ ), and the general property  $\operatorname{Re} g(z) = 0$  for  $|z| > 1$  of the reservoir correlation function. A characteristic example of  $I(\omega)$  is plotted in Fig. 1.

We observe that  $I(\omega)$  has a spectral line for  $\omega > \omega_D$ . Recalling that our result (6) only includes single-phonon transitions,<sup>10</sup> this might appear to be a contradiction. We see, however, that the line is not brought about by the transition dipole moment  $\mu_{21}$ , as usual, but by the permanent dipole moments  $\mu_{22}$  and  $\mu_{11}$  of the two levels. From the diagrams of Fig. 2 we then anticipate that it could be possible to absorb a photon with  $\omega > \omega_D$ , and in such a way that energy conservation can be met with a single-phonon process. The mismatch between  $\omega$  and  $\omega_0$  is simply the phonon frequency. Although level  $|2\rangle$  cannot be populated because of  $\omega_0 > \omega_D$ , it can nevertheless be reached via a photon-phonon process. This mechanism gives rise to the line at  $\omega > \omega_D$ .

Figure 2 suggests that the processes responsible for this line (b and c) can only occur for  $|\omega - \omega_0| < \omega_D$ , since otherwise the phonon frequency would become

larger than  $\omega_D$ . This can be shown explicitly from Eq. (6). If we work out the real part and use  $\text{Re } g(z) = 0$  for  $|z| > 1$ , we find

$$I(\omega) = 0 \quad \text{for} \quad |\omega - \omega_0| > \omega_D. \quad (7)$$

Furthermore, from Eq. (6) it follows that the line must be situated at the zero of the real part of the denominator, e.g., at the solution of  $\bar{\omega} \approx a \text{Im}(g(\hat{\bar{\omega}} - \hat{\omega}_0) + g^*(\hat{\omega}_0 - \hat{\bar{\omega}}))$ . We know that  $g(z)$  attains its extreme at  $z = \pm 1$  (for a Debye model we have  $g(\pm 1) = \pm \infty$ ) and that  $\omega > \omega_D \gg a$ , which gives for the position of the line  $\bar{\omega} = \omega_0 + \omega_D$ . Together with Eq. (7), this explains the sharp edge of the line at the blue side. In conclusion, the finite memory-time in the atom-crystal interaction prohibits the factorization of the density operator in thermal equilibrium, which in turn gives rise to a new spectral line at  $\omega_0 + \omega_D$ , provided that the system has a permanent dipole moment. Its profile is given by Eq. (6).

As a last remark we mention that the detailed shape of the line at  $\omega_0 + \omega_D$  depends crucially on the adopted model for the phonon dispersion relation. The sharp peak at  $\omega = \omega_0 + \omega_D$  is a consequence of the cutoff of the dispersion relation at  $\omega_D$ . In a more realistic model, where the detailed structure of the first Brillouin zone for a specific crystal is taken into consideration, this peak will smoothen out. Expression (6) for the line shape remains valid, but the function  $g(z)$  will behave more realistic. The infinities in  $z = \pm 1$  will turn into finite van Hove singularities.<sup>11</sup>

#### ACKNOWLEDGMENTS

This research was supported by the Office of Naval Research and the Air Force Office of Scientific Research (AFSC), United States Air Force, under Contract No. F49620-86-C-0009.


# REFERENCES


1. Z. W. Gortel, H. J. Kreuzer, P. Piercy and R. Teshima, Phys. Rev. B 27, 5066 (1983).
2. A. A. Maradudin, E. W. Montroll, G. H. Weiss and I. P. Ipatova, Theory of Lattice Dynamics in the Harmonic Approximation, Suppl. 3 of Solid State Physics (Academic, New York, 1971).
3. S. Efrima, C. Jedrzejek, K. F. Freed, E. Hood and H. Metiu, J. Chem. Phys. 79, 2436 (1983).
4. H. F. Arnoldus, S. van Smaalen and T. F. George, Adv. Chem. Phys. (1987), in press.
5. W. H. Louisell, Quantum Statistical Properties of Radiation (Wiley, New York, 1973), Ch. 6.
6. H. F. Arnoldus and T. F. George, J. Math. Phys. 28, 2731 (1987).
7. H. F. Arnoldus and T. F. George, Phys. Rev. B (1987), submitted.
8. A. Peremans, J. Darville, J. M. Gilles and T. F. George, Phys. Rev. B 35, 2690 (1987).
9. H. F. Arnoldus and T. F. George, Phys. Rev. B 36, 1 (1987).
10. A low-intensity laser can only cause transitions between two states  $|n\rangle$  and  $|m\rangle$  for which  $\langle n | H_i | m \rangle \neq 0$ , according to the Golden Rule. The Hamiltonian  $H_i$  from Eq. (1) connects only states which differ by one phonon. See for instance: S. W. Lovesey, Condensed Matter Physics, Dynamic Correlations (Benjamin, London, 1980) pg. 3.
11. N. W. Ashcroft and N. D. Mermin, Solid State Physics (Holt, Rinehart and Winston, New York, 1976) pg. 465.

# FIGURE CAPTIONS

Fig. 1. Absorption profile  $I(\omega)$  from Eq. (5) as a function of  $\hat{\omega} = \omega/\omega_D$ , and for  $\omega_0 = 3\omega_D$  (dotted line),  $a = 0.4 \times \omega_D$ ,  $\gamma = 10$  (low temperature), and  $(\mu_{22} - \mu_{11})^2 = 0.5 \times \mu_{21}^2$ . The left peak at  $\omega < \omega_D$  is the far red wing of the regression part of the line, which is brought about by transitions with  $\omega = \omega_{\text{phonon}}$  (diagram (a) of Fig. 2), and governed by the transition dipole moment  $\mu_{21}$ . The peak on the right-hand side is the memory-time-induced extra resonance, with  $|\omega - \omega_0| = \omega_{\text{phonon}}$ . Processes represented by the diagrams (b) and (c) of Fig. 2 give rise to this line, and its line shape is given by Eq. (6). In a zero memory-time approximation this line vanishes.

Fig. 2. Energy-conserving diagrams which contribute to the absorption spectrum  $I(\omega)$ . Double arrows indicate a photon and single arrows a phonon. Diagram (a) and its reverse are responsible for the profile at  $\omega < \omega_D$ , whereas diagrams (b) and (c) represent processes which yield the line at  $\omega > \omega_D$ . Since level  $|2\rangle$  is not populated, the reverse processes (stimulated emission) of diagrams (b) and (c) do not occur.





DL/1113/86/2

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
		Dr. David L. Nelson Chemistry Division Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. J. E. Jensen  
Hughes Research Laboratory  
3011 Malibu Canyon Road  
Malibu, California 90265

Dr. J. H. Weaver  
Department of Chemical Engineering  
and Materials Science  
University of Minnesota  
Minneapolis, Minnesota 55455

Dr. A. Reisman  
Microelectronics Center of North Carolina  
Research Triangle Park, North Carolina  
27709

Dr. M. Grunze  
Laboratory for Surface Science and  
Technology  
University of Maine  
Orono, Maine 04469

Dr. J. Butler  
Naval Research Laboratory  
Code 6115  
Washington D.C. 20375-5000

Dr. L. Interante  
Chemistry Department  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. Irvin Heard  
Chemistry and Physics Department  
Lincoln University  
Lincoln University, Pennsylvania 19352

Dr. K.J. Klaubunde  
Department of Chemistry  
Kansas State University  
Manhattan, Kansas 66506

Dr. C. B. Harris  
Department of Chemistry  
University of California  
Berkeley, California 94720

Dr. F. Kutzler  
Department of Chemistry  
Box 5055  
Tennessee Technological University  
Cookeville, Tennessee 38501

Dr. D. DiLella  
Chemistry Department  
George Washington University  
Washington D.C. 20052

Dr. R. Reeves  
Chemistry Department  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. Steven M. George  
Stanford University  
Department of Chemistry  
Stanford, CA 94305

Dr. Mark Johnson  
Yale University  
Department of Chemistry  
New Haven, CT 06511-8118

Dr. W. Knauer  
Hughes Research Laboratory  
3011 Malibu Canyon Road  
Malibu, California 90265



ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. G. A. Somorjai  
Department of Chemistry  
University of California  
Berkeley, California 94720

Dr. J. Murday  
Naval Research Laboratory  
Code 6170  
Washington, D.C. 20375-5000

Dr. J. B. Hudson  
Materials Division  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. Theodore E. Madey  
Surface Chemistry Section  
Department of Commerce  
National Bureau of Standards  
Washington, D.C. 20234

Dr. J. E. Demuth  
IBM Corporation  
Thomas J. Watson Research Center  
P.O. Box 218  
Yorktown Heights, New York 10598

Dr. M. G. Lagally  
Department of Metallurgical  
and Mining Engineering  
University of Wisconsin  
Madison, Wisconsin 53706

Dr. R. P. Van Duyne  
Chemistry Department  
Northwestern University  
Evanston, Illinois 60637

Dr. J. M. White  
Department of Chemistry  
University of Texas  
Austin, Texas 78712

Dr. D. E. Harrison  
Department of Physics  
Naval Postgraduate School  
Monterey, California 93940

Dr. R. L. Park  
Director, Center of Materials  
Research  
University of Maryland  
College Park, Maryland 20742

Dr. W. T. Peria  
Electrical Engineering Department  
University of Minnesota  
Minneapolis, Minnesota 55455

Dr. Keith H. Johnson  
Department of Metallurgy and  
Materials Science  
Massachusetts Institute of Technology  
Cambridge, Massachusetts 02139

Dr. S. Sibener  
Department of Chemistry  
James Franck Institute  
5640 Ellis Avenue  
Chicago, Illinois 60637

Dr. <sup>Arnold</sup> Arnold Green  
Quantum Surface Dynamics Branch  
Code 3817  
Naval Weapons Center  
China Lake, California 93555

Dr. A. Wold  
Department of Chemistry  
Brown University  
Providence, Rhode Island 02912

Dr. S. L. Bernasek  
Department of Chemistry  
Princeton University  
Princeton, New Jersey 08544

Dr. W. Kohn  
Department of Physics  
University of California, San Diego  
La Jolla, California 92037

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. F. Carter  
Code 6170  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. Richard Colton  
Code 6170  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. Dan Pierce  
National Bureau of Standards  
Optical Physics Division  
Washington, D.C. 20234

Dr. R. Stanley Williams  
Department of Chemistry  
University of California  
Los Angeles, California 90024

Dr. R. P. Messmer  
Materials Characterization Lab.  
General Electric Company  
Schenectady, New York 22217

Dr. Robert Gomer  
Department of Chemistry  
James Franck Institute  
5640 Ellis Avenue  
Chicago, Illinois 60637

Dr. Ronald Lee  
R301  
Naval Surface Weapons Center  
White Oak  
Silver Spring, Maryland 20910

Dr. Paul Schoen  
Code 6190  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. John T. Yates  
Department of Chemistry  
University of Pittsburgh  
Pittsburgh, Pennsylvania 15260

Dr. Richard Greene  
Code 5230  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. L. Kesmodel  
Department of Physics  
Indiana University  
Bloomington, Indiana 47403

Dr. K. C. Janda  
University of Pittsburgh  
Chemistry Building  
Pittsburg, PA 15260

Dr. E. A. Irene  
Department of Chemistry  
University of North Carolina  
Chapel Hill, North Carolina 27514

Dr. Adam Heller  
Bell Laboratories  
Murray Hill, New Jersey 07974

Dr. Martin Fleischmann  
Department of Chemistry  
University of Southampton  
Southampton SO9 5NH  
UNITED KINGDOM

Dr. H. Tachikawa  
Chemistry Department  
Jackson State University  
Jackson, Mississippi 39217

Dr. John W. Wilkins  
Cornell University  
Laboratory of Atomic and  
Solid State Physics  
Ithaca, New York 14853

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. R. G. Wallis  
Department of Physics  
University of California  
Irvine, California 92664

Dr. D. Ramaker  
Chemistry Department  
George Washington University  
Washington, D.C. 20052

Dr. J. C. Hemminger  
Chemistry Department  
University of California  
Irvine, California 92717

Dr. T. F. George  
Chemistry Department  
University of Rochester  
Rochester, New York 14627

Dr. G. Rubloff  
IBM  
Thomas J. Watson Research Center  
P.O. Box 218  
Yorktown Heights, New York 10598

Dr. Horia Metiu  
Chemistry Department  
University of California  
Santa Barbara, California 93106

Dr. W. Goddard  
Department of Chemistry and Chemical  
Engineering  
California Institute of Technology  
Pasadena, California 91125

Dr. P. Hansma  
Department of Physics  
University of California  
Santa Barbara, California 93106

Dr. J. Baldeschwieler  
Department of Chemistry and  
Chemical Engineering  
California Institute of Technology  
Pasadena, California 91125

Dr. J. T. Keiser  
Department of Chemistry  
University of Richmond  
Richmond, Virginia 23173

Dr. R. W. Plummer  
Department of Physics  
University of Pennsylvania  
Philadelphia, Pennsylvania 19104

Dr. E. Yeager  
Department of Chemistry  
Case Western Reserve University  
Cleveland, Ohio 44106

Dr. N. Winograd  
Department of Chemistry  
Pennsylvania State University  
University Park, Pennsylvania 16802

Dr. Roald Hoffmann  
Department of Chemistry  
Cornell University  
Ithaca, New York 14853

Dr. A. Steckl  
Department of Electrical and  
Systems Engineering  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. G.H. Morrison  
Department of Chemistry  
Cornell University  
Ithaca, New York 14853

END

DATE

FILMED

6-1988

DTIC